INDEX

A.I.V.-fodder, 144–52.
Ammonium phosphate, for rice in Ceylon, 124–5.
Ammonium sulphate, for cotton in Sudan, 277, 280–4, 293–4.
— for rice in Australia, 233.
— for rice in Ceylon, 124.
— for tea in Ceylon, 107–8.
— trials with, on arable crops, 129–43.
Arnold, C. W. B., Nyasaland tobacco industry, 379–83.
Australia, rice-growing in, 229–36.

Bal, D. V., black cotton soils of Central Provinces, India, 261–8.
Barley, as forage crop in W. Canada, 326.
— effect of N-fertilizers on, 130–5, 137.
Bartel, L. H., *see* Rhodes, C. C.
Basic slag, for acidic upland soils, 51 sqq.
Beckley, V. A., chlorosis and die-back in coffee, 203–9.
Beef, market for, in Gt. Britain, 160–73; *per capita* consumption, 162–3, 167, 171–2; prices, 164–72; supplies, 162–4, 168–70.
Blood-meal, as poultry-feed, 216 sqq.
— for tea in Ceylon, 107–8.
British Guiana, rice industry of, 237–47.

Calcium cyanamide, effect on charlock in barley, 46–9; on germinating barley, 41–6.
— for tea in Ceylon, 107–8.
— trials with, on arable crops, 129–43.
Callow, E. H., carcass quality of the pig in relation to growth and diet, 80–104.
Canada, Western, forage-crop production in dry areas, 320–30; major soil zones in prairie provinces of, 320–1; grassland management, 328–9; hay and pasture crops, 322–6; seed-production of grasses and legumes, 327–8.
Cereals, as forage crops in W. Canada, 322.
Ceylon, climate and soils, 119–20; cultivation of rice, 119–28.
Charlock, in barley, effect of cyanamide on, 46–9.
Chippindale, H. G., *see* Davies, R. O.
Clover, red, effects of cutting and grazing on dry-matter content of, 197–202.
— sweet, in W. Canada, 324–5, 327.
— *Trifolium Johnstonii*, 155–9.
Clover, wild white, effect of grazing on dry-matter content of, 199.
— effect of pH of soil and soil-drainage, 54, 55, 56.
— protein-synthesis by, 57–8.
— response of, to treatment on acidic soils, 50–9.
Coffee (*Coffea arabica*), observations on, in Kenya, 203–14; cherries, effect of chlorosis on, 205; chlorosis and die-back in, 203-9; root-systems, 206, 210; vegetative propagation of, 210-14.
Cotton, American, effect of shade on, 31–40.
— irrigated, in Sudan, experiments on factors controlling production of, 276–88, 289–94.
— — effect on yield of N-response, 282–3, 293; of sowing-date, 281–2, 287, 291–2; of spacing, 282, 293–4; interactions of these factors, 283–7, 288, 290–1.
— — Gezira irrigation scheme, 276 sqq.
— — pests, 278–9, 289–91, 293.
— — quality of cotton, 293–4.
— — rainfall, 278, 279, 292.
— — soils, 277–8, 278–9.
Cows, beef from, 172.
— feeding-trial with, 146–51.
— grass tetany in, 371–3.
— milk fever in, 365–6.
Crampton, E. W., effect of fertilization on nutritive value of pasture grass, 331–45.
Cropping, mixed, in primitive agriculture, 189–95.
Crowther, E. M., comparative trials of cyanamide and other nitrogenous fertilizers on arable crops, 129–43.
Crowther, F., *see* Lambert, A. R.

Dash, J. Sydney, rice industry in British Guiana, 237–47.
Davies, R. O., response to treatment of grasses and clover on acidic upland soils, and effect of herbage plants on reaction of acidic soils, 50–9.
Dicyanodiamide, effect of, on N-loss in soils, 142–3.
Diseases, of coffee, 203–9.
— mineral deficiency, of larger domesticated animals, 363–78; aphosphorosis, 364, 368–9; fluorosis, 376; goitre, 375–6; hypocalcaemia, 365–6, 372–3; hypomagnesaemia, 371–3; nutritional anaemias, 373–5; osteofibrosis, 369; 'salt-hunger', 376–7; skeletal disorders, 366.
— of rice in Ceylon, 128.
'Doppellender' calf, carcass of, 3.

Eden, T., studies in the yield of tea, 104–18.
Edwards, D. C., grasslands of Kenya, 153–9.
Eggs, hen's, effects of protein diets on weight, size, and number, 215–28, 313–19; short-period *v.* yearly recording, 316–17.
— — factors influencing production of, 13–20.

Evans, D. I., *see* Eden, T.
Evans, H., root-system of the sugar-cane, I, methods of study, 351–62.

Farming, dry-land, in prairie provinces of W. Canada, 320–30.
Fertilizers, effect of, on composition of made tea, 115–18.
— effect of, on N-content of soil drainage water, 54–6.
— effect of, on herbage plants, &c., 50 sqq.
— effect of, on nutritive value of pasture grass, 331–45.
— for rice in Australia, 233.
— for rice in British Guiana, 243–4.
— for rice in Ceylon, 124–5.
— for tea in Ceylon, 113–14.
— nitrogenous, for coffee, 205.
— — trials with, on arable crops, 129–43.
— use of, in expts. on grasses and clover, 50 sqq.
Finlayson, D. A., *see* Crampton, E. W.
Forage crops, in prairie provinces of W. Canada, 320–30.
Fraser, A. H. H., wool-growth in sheep as affected by carbohydrate in diet, II, 75–9.

Gillett, S., vegetative propagation of coffee, 210–14.
Godden, W., *see* Watson, C. J.
Grass, artificially dried, as food for sheep and rabbits, 346–50.
— Kikuyu, in Kenya, 155–9.
— silage, feeding-value of, 144–52.
— *Themeda triandra* Forsk., 153–9.
Grasses and clover, dry-matter content of, 197–202.
— in dry-land agric. in W. Canada, 322–4, 326.
— on acidic upland soils, response to treatment of, 50–9.
Grassland management, in Kenya, 157–8.
— on ranges in W. Canada, 328–9.
Grasslands of Kenya, 153–9.
Great Britain, development of market for beef in, 160–73.
Green, H. H., minerals in relation to disease of larger domesticated animals, 363–78.
Green-manuring, for tea in Ceylon, 124.
Greenwood, A. W., inheritance of productivity in farm live stock: breeding for egg-production, 13–20.

Hammond, J., inheritance of productivity in farm live stock: meat, 1–6.
Hanley, F., *see* Nicholson, H. H.
Hay crops in W. Canada, 322–6.
Horse, evolution in skeleton of, 1–2.
— osteofibrosis in, 369–71; transit tetany in, 373.

India, black cotton soils of Central Provinces, 261–8.
Irrigation, rice in Australia, 229–31, 234; in British Guiana, 238–9; in Ceylon, 120, 125–6.
— on ranges in W. Canada, 329.

Jones, H. Trefor, soils of Yorkshire, 269–75.
Jooste, P. E. F., *see* Rhodes, C. C.

Kenya, coffee in, 203–14.
— grasslands of, 153–9.
Kirk, L. E., forage-crop production in dry-land agriculture and on ranges in W. Canada, 320–30.
Knight, R. L., effect of shade on American cotton, 31–40.

Lambert, A. R., experiments on interrelation of factors controlling cotton-production in Sudan, I, 276–88; II, 289–94.
Legumes in mixed cropping, 189–95.
Live stock, inheritance of productivity in, 1–30.
— mineral deficiency diseases of, 363–78.
Lord, L., cultivation of rice in Ceylon, 119–28.
Lucerne (alfalfa), in W. Canada, 323–4, 326–7, 329.
— meal for rabbits, 333–5, 343.
Lush, J. L., inheritance of productivity in live stock: discussion, 25–30.

Maize, as forage crop in W. Canada, 322.
Maize-meal, as food for poultry, 216 sqq.
Maize-starch, effect on wool-growth in sheep, 75–9.
Meat supplies in Gt. Britain, 160–1, 168.
Milk, analyses of, from different sources, 86.
— from cows, as food for pigs, 87.
— — fed on A.I.V.-fodder, 148–50.
— — genetic and other influences on yield of, 7–12.
Milton, W. E. J., dry-matter content of certain grass and clover species, 197–202.
Minerals in relation to disease of larger domesticated animals, 363–78.
Murray, K. A. H., development of market for beef in Great Britain, 160–73.

Nichols, J. E., inheritance of productivity in live stock: wool, 21–4.
— *see* Fraser, A. H. H.
Nicholson, H. H., soil conditions in East Anglia, 60–74.
Nicol, Hugh, mixed cropping in primitive agriculture, 189–95.
Nitro-chalk, for acidic upland soils, 51 sqq.
Nitrogen, content of, in made tea, 117; in tea-plant, 112.
— effect of manurial, on made tea, 117–18.

Nitrogen, effect of manurial, on coffee with die-back, 205.
— in soil-sand mixture, effect of grasses and clover on, 57–8.
— loss of, by leaching, 142–3.
— nitrification of, in ammonia, 266–7; in oil-cake, 266.
— organic and inorganic, for tea, 108.
— removal of, from tea soil, 112–13.
— removed from soil by grasses, 56–7.
— response of tea to, 107–8.
Nyasaland, tobacco industry in, 379–83.

Oats, as forage crop in W. Canada, 326.
— winter, effect of N-fertilizers on, 137–9.
Ogg, W. G., soils of Scotland, I, 174–88; II, 248–60; III, 295–312.

Pasture crops in W. Canada, 322–6.
Pasture grass, effect of fertilizers on nutritive value of, 331–2, 336–42, 344–5; on quality of protein in, 338–40, 345.
Pasture herbage, digestibility of artificially dried, by sheep and rabbits, 346–50.
Patterson, J. B. E., comparative feeding-values of A.I.V.-grass silage and a ration containing mangolds and hay, 144–52.
Peat in Scotland, 181–4.
Pests, of cotton, irrigated, in Sudan, 36–8, 278, 279, 281, 288, 289–91.
— of rice in British Guiana, 244; in Ceylon, 127–8.
Phosphoric acid, content of, in tea-plant, 112.
— in Ceylon soils, 124–5.
— in soils of NE. Scotland, 260.
— removal of, from tea soil, 113.
Pigs, breeds of, for pork and bacon, 96.
— carcass quality of, in relation to growth and diet, 80–104.
— — common defects in, 96–7.
— — for pork and bacon, 96, 100.
— — ideal, 95.
— — measurement of, 95–7.
— deficiency diseases of, 87–90, 369, 371, 374–5.
— diet, effect of, on growth and conformation, 84 sqq.; amino acids in, 89; fat-content of, 91; protein-deficiency in, 89; use of cow's milk and green foods in, 87.
— fat, composition and properties of, 90–4.
— — effect of age and diet on hardness of, 94.
— — — diet on, 92–3.
— — — dietetic deficiencies on, 94.
— — — external temperature on, 93.
— — — fish-oils on, 98–9.
— — quality of, 97–100.
— — rancidity in, 97–9.
— genetic characters of, for bacon, pork, and lard, 4–6.
— milk, composition of, 86.
Pigs, muscular tissue (and bone), effect of diet on, 94–5.
— — electrical resistance of, 100–1.
— time taken to mature, 81–2; growth-rate and length, 83–4.
Plants, herbage, effect of, on soil pH and drainage, 52–4.
Potash, content of, in tea-plant, 112.
— for acidic upland soils, 51 sqq.
— in pastures, 376–7.
— in soils of NE. Scotland, 260.
— removal of, from tea soil, 113.
— response of tea to, 107.
Productivity, inheritance of, in cattle 2–3; in dairy cows 7–12; in hens 13–20; in live stock 25–30; in pigs 4–6; in sheep 3, 21–4.
Protein, dietary, effect on White Leghorn pullets, 215–28, 313–19.
— effect of adding to rabbit-feed, 332–6.
— in fresh grass and A.I.V.-fodder, 145–6.
— in grasses, &c., synthesis of, 57–8.
— in pasture grass, effect of fertilizers on quality of, 338–40, 345.

Rabbits, use of, in nutritional experiments, 333–45, 346–50.
Reed canary-grass, nutritive value of, 332–6, 343.
Rhodes, C. C., effects of varying amounts of animal protein fed to White Leghorn pullets, I, 215–28, II, 313–19.
Rice, in Australia, 229–36; crop rotations, 232–3; cultural operations, 230–1; fertilizers, 233; marketing and economics, 235–6; Murrumbidgee irrigation area, 228–30, 234, 236; research, 234; statistics, 229; varieties, 233–4.
— in British Guiana, 237–47; cultivation, 237–40; fertilizers, 243–4; irrigation, 238–9; marketing and economics, 244–7; pests, 244; research, 240–4; statistics, 238–40, 246; varieties, 243.
— in Ceylon; cultivation of, 112–13, 126; irrigation, 120, 125–6; manuring, 123–5; pests and diseases, 127–8; varieties, 121–2; yields, 119, 126–7.
Richardson, H. L., action of cyanamide on germinating seeds and on charlock in barley, 41–9.
Root crops, effect of N-fertilizers on, 134, 136–7.
Rye, as forage crop in W. Canada, 326.

Scotland, agriculture, 176; climate, 174; geology, 174–5; peat, 181–4; soils, 178–80.
— Central Valley and Southern Uplands, 295–312; climate and geology, 295–6; moorland, 296–7; soils, 297–312; vegetation, 296–7.

Scotland, Highlands and Hebrides, 180–8; peat, 181–4; rocks, 180; soils, 184–8; vegetation, 180–1.
— NE. region, 248–60; climate and geology, 248; soils, 249–60; vegetation, 248–9.
Seed-production of grasses and legumes in prairie provinces of W. Canada, 326–8.
Sheep, genetic characters of, for mutton and lamb, 3.
— — for wool, 21–4.
— fluorosis in, 376.
— goitre in, 375–6.
— wool-growth, as affected by dietetic carbohydrate, 75–9.
Smith, A. D. Buchanan, inheritance of productivity in farm live stock: milk, 7–12.
Soil conditions in East Anglia, 60–74.
— drainage, at Craibstone, 259.
— — nitrate-N concentration in, 54–6.
— fertility estimations, NE. Scotland, 259–60.
— *Molinia*, effect of herbage plants on, 50–9.
— nitrogen, as affected by grasses, 56–8.
— nitrate, as affected by winter leaching, 142–3.
— pH of, from grass plots, 52–4.
— — cultivated, in NE. Scotland, 259–60.
— uncropped, effect of manures on nitrate-N in drainage from, 55.
Soils, acidic, effect of herbage plants on reaction of, 50–9.
— alkali, in W. Canada, crops for, 325–6.
— black cotton, in Central Provinces, India, 261–8; biological aspects, 264–7; composition, 261–4; moisture-content, 265; nitrification, 265–7; organic matter, 265.
— cotton, in Sudan, 277–8, 278–9.
— in Ceylon, cultivation of, 119–28.
— in East Anglia, 60–74; alluvial, 66–9; Boulder Clay, 62–5; Chalk, 69–71; Clay and Gault, 71–4; Glacial Gravel, 65–6; Lower Greensand, 74.
— of Scotland, 174–88, 248–60, 295–312; brown, 178, 180, 257–9, 307–11; gley, 178, 186–7, 256–7, 306–7; gley-podsolic, 178, 185–7, 254–6, 303–6; podsolic, 178, 184–5, 249–54, 298–302; with undeveloped profile, 178, 180, 187–8, 311–12.
Soils of Yorkshire, 269–75; climate and geology, 269–71; podsolic, 271–3; post-glacial, 273–5.
— rice, in Ceylon, 120; irrigation of, 120, 125–6.
Sudan, Anglo-Egyptian, experiments on irrigated cotton in, 276–88, 289–94.
Sugar-cane, root-system of, 351–62.

Tea, composition of made, 117–18.
— determination of constituents of made, 116–17.
— jats, 110–11.
— quality of, 115.
— response of, to nitrogen and potash, 105–12.
— studies in yield of, 105–18.
— yields of leaf and wood, 111–12.
Timothy grass, dry-matter content of, 198–9.
— nutritive value of, 332–6, 343.
— protein in, 145.
Tobacco, in Nyasaland, 379–83; air-cured and sun-cured, 383; dark fired, 381–2; flue-cured, 382–3.
Trench, A. D., *see* Beckley, V. A.

Vegetative propagation of coffee, 210–14.
Vitamin D in animal nutrition, 368–9.
Vitamins A and D in pigs' dietary, 87, 98–9.

Watson, C. J., comparative digestibility of artificially-dried pasture herbage by sheep and rabbits, 346–50.
Watson, J. A. Scott, *see* Murray, K. A. H.
West, E. S., rice-growing in Australia, 229–36.
Wheat, winter, effect of N-fertilizers on, 139–42.
White Leghorn pullets, effects of protein-dietary on, 215–28, 313–19.
Wool, sheep's, effect of dietetic carbohydrate on fibre-length, 77–8; fineness, 78–9; fleece-weight, 76–7.
— inheritance of fleece-characters, 22–4.

ERRATA

Vol. III, July 1935, p. 278, line 11 from top, *in lieu of* Dolba Grange *read* Dolga Grange; p. 284, line 21 from top, *in lieu of* 0·75 *read* 0·73; p. 288, line 19 from top, *in lieu of* sown *read* sowing-date.

PRINTED IN GREAT BRITAIN AT THE UNIVERSITY PRESS, OXFORD
BY JOHN JOHNSON, PRINTER TO THE UNIVERSITY